

ON TREE-PARTITION-WIDTH

DAVID R. WOOD

ABSTRACT. A *tree-partition* of a graph G is a proper partition of its vertex set into ‘bags’, such that identifying the vertices in each bag produces a forest. The *tree-partition-width* of G is the minimum number of vertices in a bag in a tree-partition of G . An anonymous referee of the paper by Ding and Oporowski [*J. Graph Theory*, 1995] proved that every graph with tree-width $k \geq 3$ and maximum degree $\Delta \geq 1$ has tree-partition-width at most $24k\Delta$. We prove that this bound is within a constant factor of optimal. In particular, for all $k \geq 3$ and for all sufficiently large Δ , we construct a graph with tree-width k , maximum degree Δ , and tree-partition-width at least $(\frac{1}{8} - \epsilon)k\Delta$. Moreover, we slightly improve the upper bound to $\frac{5}{2}(k+1)(\frac{7}{2}\Delta - 1)$ without the restriction that $k \geq 3$.

1. INTRODUCTION

A graph¹ H is a *partition* of a graph G if:

- each vertex of H is a set of vertices of G (called a *bag*),
- every vertex of G is in exactly one bag of H , and
- distinct bags A and B are adjacent in H if and only if some edge of G has one endpoint in A and the other endpoint in B .

The *width* of a partition is the maximum number of vertices in a bag. Informally speaking, the graph H is obtained from a proper partition of $V(G)$ by identifying the vertices in each part, deleting loops, and replacing parallel edges by a single edge.

If a forest T is a partition of a graph G , then T is a *tree-partition* of G . The *tree-partition-width*² of G , denoted by $\text{tpw}(G)$, is the minimum width of a tree-partition of G . Tree-partitions were independently introduced by Seese [23] and Halin [19], and have since been widely investigated [6, 7, 12, 13, 17, 24]. Applications of tree-partitions include graph drawing [9, 14, 15, 25], graph colouring [2], partitioning graphs into subgraphs with only small components [1], monadic second-order logic [20], and network emulations [3, 4, 8, 18]. Planar-partitions and other more general structures have also recently been studied [11, 25].

Date: February 2, 2008.

1991 *Mathematics Subject Classification.* 05C70 (graph theory: factorization, matching, covering and packing).

Key words and phrases. graph, tree-partition, tree-partition-width, tree-width.

Research supported by a Marie Curie Fellowship of the European Community under contract MEIF-CT-2006-023865, and by the projects MEC MTM2006-01267 and DURSI 2005SGR00692.

¹All graphs considered are undirected, simple, and finite. Let $V(G)$ and $E(G)$ respectively be the vertex set and edge set of a graph G . Let $\Delta(G)$ be the maximum degree of G .

²Tree-partition-width has also been called *strong tree-width* [7, 23].

What bounds can be proved on the tree-partition-width of a graph? Let $\text{tw}(G)$ denote the tree-width³ of a graph G . Seese [23] proved the lower bound,

$$2 \text{tpw}(G) \geq \text{tw}(G) + 1.$$

In general, tree-partition-width is not bounded from above by any function solely of tree-width. For example, wheel graphs have bounded tree-width and unbounded tree-partition-width [7]. However, tree-partition-width is bounded for graphs of bounded tree-width *and* bounded degree [12, 13]. The best known upper bound is due to an anonymous referee of the paper by Ding and Oporowski [12], who proved that

$$\text{tpw}(G) \leq 24 \text{tw}(G) \Delta(G)$$

whenever $\text{tw}(G) \geq 3$ and $\Delta(G) \geq 1$. Using a similar proof, we make the following improvement to this bound without the restriction that $\text{tw}(G) \geq 3$.

Theorem 1. *Every graph G with tree-width $\text{tw}(G) \geq 1$ and maximum degree $\Delta(G) \geq 1$ has tree-partition-width*

$$\text{tpw}(G) < \frac{5}{2}(\text{tw}(G) + 1) \left(\frac{7}{2} \Delta(G) - 1 \right).$$

Theorem 1 is proved in Section 2. Note that Theorem 1 can be improved in the case of chordal graphs. In particular, a simple extension of a result by Dujmović et al. [14] implies that

$$\text{tpw}(G) \leq \text{tw}(G)(\Delta(G) - 1)$$

for every chordal graph G with $\Delta(G) \geq 2$; see [24] for a simple proof. Nevertheless, the following theorem proves that $\mathcal{O}(\text{tw}(G) \Delta(G))$ is the best possible upper bound, even for chordal graphs.

Theorem 2. *For every $\epsilon > 0$ and integer $k \geq 3$, for every sufficiently large integer $\Delta \geq \Delta(k, \epsilon)$, for infinitely many values of N , there is a chordal graph G with N vertices, tree-width $\text{tw}(G) \leq k$, maximum degree $\Delta(G) \leq \Delta$, and tree-partition-width*

$$\text{tpw}(G) \geq \left(\frac{1}{8} - \epsilon \right) \text{tw}(G) \Delta(G).$$

Theorem 2 is proved in Section 3. Note that Theorem 2 is for $k \geq 3$. For $k = 1$, every tree is a tree-partition of itself with width 1. For $k = 2$, we prove that the upper bound $\mathcal{O}(\Delta(G))$ is again best possible; see Section 4.

2. UPPER BOUND

In this section we prove Theorem 1. The proof relies on the following separator lemma by Robertson and Seymour [22].

Lemma 1 ([22]). *For every graph G with tree-width at most k , for every set $S \subseteq V(G)$, there are edge-disjoint subgraphs G_1 and G_2 of G such that $G_1 \cup G_2 = G$, $|V(G_1) \cap V(G_2)| \leq k + 1$, and $|S - V(G_i)| \leq \frac{2}{3}|S - (V(G_1) \cap V(G_2))|$ for each $i \in \{1, 2\}$.*

³A graph is *chordal* if every induced cycle is a triangle. The *tree-width* of a graph G can be defined to be the minimum integer k such that G is a subgraph of a chordal graph with no clique on $k + 2$ vertices. This parameter is particularly important in algorithmic and structural graph theory; see [5, 21] for surveys.

Theorem 1 is a corollary of the following stronger result.

Lemma 2. *Let $\alpha := 1 + 1/\sqrt{2}$ and $\gamma := 1 + \sqrt{2}$. Let G be a graph with tree-width at most $k \geq 1$ and maximum degree at most $\Delta \geq 1$. Then G has tree-partition-width*

$$\text{tpw}(G) \leq \gamma(k+1)(3\gamma\Delta - 1) .$$

Moreover, for each set $S \subseteq V(G)$ such that

$$(\gamma+1)(k+1) \leq |S| \leq 3(\gamma+1)(k+1)\Delta,$$

there is a tree-partition of G with width at most

$$\gamma(k+1)(3\gamma\Delta - 1),$$

such that S is contained in a single bag containing at most $\alpha|S| - \gamma(k+1)$ vertices.

Proof. We proceed by induction on $|V(G)|$.

Case 1. $|V(G)| < (\gamma+1)(k+1)$: Then no set S is specified, and the tree-partition in which all the vertices are in a single bag satisfies the lemma. Now assume that $|V(G)| \geq (\gamma+1)(k+1)$, and without loss of generality, S is specified.

Case 2. $|V(G) - S| < (\gamma+1)(k+1)$: Then the tree-partition in which S is one bag and $V(G) - S$ is another bag satisfies the lemma. Now assume that $|V(G) - S| \geq (\gamma+1)(k+1)$.

Case 3. $|S| \leq 3(\gamma+1)(k+1)$: Let N be the set of vertices in G that are adjacent to some vertex in S but are not in S . Then $|N| \leq \Delta|S| \leq 3(\gamma+1)(k+1)\Delta$. If $|N| < (\gamma+1)(k+1)$ then add arbitrary vertices from $V(G) - (S \cup N)$ to N until $|N| \geq (\gamma+1)(k+1)$. This is possible since $|V(G) - S| \geq (\gamma+1)(k+1)$.

By induction, there is a tree-partition of $G - S$ with width at most $\gamma(k+1)(3\gamma\Delta - 1)$, such that N is contained in a single bag. Create a new bag only containing S . Since all the neighbours of S are in a single bag, we obtain a tree-partition of G . (S corresponds to a leaf in the pattern.) Since $|S| \geq (\gamma+1)(k+1)$, it follows that $|S| \leq \alpha|S| - \gamma(k+1)$ as desired. Now $|S| \leq 3(\gamma+1)(k+1) < \gamma(k+1)(3\gamma\Delta - 1)$. Since the other bags do not change we have the desired tree-partition of G .

Case 4. $|S| \geq 3(\gamma+1)(k+1)$: By Lemma 1, there are edge-disjoint subgraphs G_1 and G_2 of G such that $G_1 \cup G_2 = G$, $|V(G_1) \cap V(G_2)| \leq k+1$, and $|S - V(G_i)| \leq \frac{2}{3}|S - (V(G_1) \cap V(G_2))|$ for each $i \in \{1, 2\}$. Let $Y := V(G_1) \cap V(G_2)$. Let $a := |S \cap Y|$ and $b := |Y - S|$. Thus $a + b \leq k+1$. Let $p_i := |(S \cap V(G_i)) - Y|$. Then $p_1 \leq 2p_2$ and $p_2 \leq 2p_1$. Let $S_i := (S \cap V(G_i)) \cup Y$. Note that $|S_i| = p_i + a + b$.

Now $p_1 + p_2 + a = |S| \geq 3(\gamma+1)(k+1)$. Thus $3p_i + a \geq 3(\gamma+1)(k+1)$ and $3p_i + 3a + 3b \geq 3(\gamma+1)(k+1)$. That is, $|S_i| \geq (\gamma+1)(k+1)$ for each $i \in \{1, 2\}$.

Now $p_1 + p_2 + a \leq 3(\gamma+1)(k+1)\Delta$. Thus $\frac{3}{2}p_i + a \leq 3(\gamma+1)(k+1)\Delta$ and $p_i \leq 2(\gamma+1)(k+1)\Delta$. Thus $p_i + a + b \leq 2(\gamma+1)(k+1)\Delta + (k+1)$. Hence $|S_i| = p_i + a + b < 3(\gamma+1)(k+1)\Delta$.

Thus we can apply induction to the set S_i in the graph G_i for each $i \in \{1, 2\}$. We obtain a tree-partition of G_i with width at most $\gamma(k+1)(3\gamma\Delta - 1)$, such that S_i is contained in a single bag T_i containing at most $\alpha|S_i| - \gamma(k+1)$ vertices.

Construct a partition of G by uniting T_1 and T_2 . Each vertex of G is in exactly one bag since $V(G_1) \cap V(G_2) = Y \subseteq S_i \subseteq T_i$. Since G_1 and G_2 are edge-disjoint, the pattern of




FIGURE 1. Illustration of Case 4.

this partition of G is obtained by identifying one vertex of the pattern of the tree-partition of G_1 with one vertex of the pattern of the tree-partition of G_2 . Since the patterns of the tree-partitions of G_1 and G_2 are forests, the pattern of the partition of G is a forest, and we have a tree-partition of G .

Moreover, S is contained in a single bag $T_1 \cup T_2$ and

$$\begin{aligned}
 |T_1 \cup T_2| &= |T_1| + |T_2| - |Y| \\
 &\leq \alpha|S_1| - \gamma(k+1) + \alpha|S_2| - \gamma(k+1) - (a+b) \\
 &= \alpha(p_1 + a + b) - \gamma(k+1) + \alpha(p_2 + a + b) - \gamma(k+1) - (a+b) \\
 &= \alpha(p_1 + p_2 + a) - 2\gamma(k+1) + (\alpha-1)a + (2\alpha-1)b \\
 &\leq \alpha|S| - 2\gamma(k+1) + (2\alpha-1)(a+b) \\
 &\leq \alpha|S| - 2\gamma(k+1) + (2\alpha-1)(k+1) \\
 &= \alpha|S| - \gamma(k+1) .
 \end{aligned}$$

Thus $|T_1 \cup T_2| \leq \alpha \cdot 3(\gamma+1)(k+1)\Delta - \gamma(k+1) = \gamma(k+1)(3\gamma\Delta - 1)$. Since the other bags do not change we have the desired tree-partition of G . \square

3. GENERAL LOWER BOUND

The remainder of the paper studies lower bounds on the tree-partition-width. The graphs employed are chordal. We first show that tree-partitions of chordal graphs can be assumed to have certain useful properties.

Lemma 3. *Every chordal graph G has a tree-partition T with width $\text{tpw}(G)$, such that for every independent set S of simplicial⁴ vertices of G , and for every bag B of T , either $B = \{v\}$ for some vertex $v \in S$, or the induced subgraph $G[B - S]$ is connected.*

⁴A vertex is *simplicial* if its neighbourhood is a clique.

Proof. Let T_0 be a tree-partition of a chordal graph G with width $\text{tpw}(G)$. Let T be the partition of G obtained from T_0 by replacing each bag B of T_0 by bags corresponding to the connected components of $G[B]$. Then T has width at most $\text{tpw}(G)$.

To prove that T is a forest, suppose on the contrary that T contains an induced cycle C . Since each bag in C induces a connected subgraph of G , G contains an induced cycle D with at least one vertex from each bag in C . Since G is chordal, D is a triangle. Thus C is a triangle, implying that the vertices in D were in distinct bags in T_0 (since the bags of T that replaced each bag of T_0 form an independent set). Hence the bags of T_0 that contain D induce a triangle in T_0 , which is the desired contradiction since T_0 is a forest. Hence T is a forest.

Let S be an independent set of simplicial vertices of G . Consider a bag B of T . By construction, $G[B]$ is connected. First suppose that $B \subseteq S$. Since S is an independent set and $G[B]$ is connected, $B = \{v\}$ for some vertex $v \in S$.

Now assume that $B - S \neq \emptyset$. Suppose on the contrary that $G[B - S]$ is disconnected. Thus $B \cap S$ is a cut-set in $G[B]$. Let v and w be vertices in distinct components of $G[B - S]$ such that the distance between v and w in $G[B]$ is minimised. (This is well-defined since $G[B]$ is connected.) Since S is an independent set, every shortest path between v and w in $G[B]$ has only two edges. That is, v and w have a common neighbour x in $B \cap S$. Since x is simplicial, v and w are adjacent. This contradiction proves that $G[B - S]$ is connected. \square

The next lemma is the key component of the proof of Theorem 2. For integers $a < b$, let $[a, b] := \{a, a + 1, \dots, b\}$ and $[b] := [1, b]$.


Lemma 4. *For all integers $k \geq 2$ and $\Delta \geq 3k + 1$, for infinitely many values of N there is a chordal graph G with N vertices, tree-width $\text{tw}(G) = 2k - 1$, maximum degree $\Delta(G) \leq \Delta$, and tree-partition-width $\text{tpw}(G) > \frac{1}{4}k(\Delta - 3k)$.*

Proof. Let n be an integer with $n > \max\{\frac{1}{2}k(\Delta - 3k), 2\}$. Let H be the graph with vertex set $\{(x, y) : x \in [n], y \in [k]\}$, where distinct vertices (x_1, y_1) and (x_2, y_2) are adjacent if and only if $|x_1 - x_2| \leq 1$. The set of vertices $\{(x, y) : y \in [k]\}$ is the x -column. The set of vertices $\{(x, y) : x \in [n]\}$ is the y -row. Observe that each column induces a k -vertex clique, and each row induces an n -vertex path.

Let C be an induced cycle in H . If (x, y) is a vertex in C with x minimum then the two neighbours of (x, y) in C are adjacent. Thus C is a triangle. Hence H is chordal. Observe that each pair of consecutive columns form a maximum clique of $2k$ vertices in H . Thus H has tree-width $2k - 1$. Also note that H has maximum degree $3k - 1$.

An edge of H between vertices (x, y) and $(x + 1, y)$ is *horizontal*. As illustrated in Figure 2, construct a graph G from H as follows. For each horizontal edge vw of H , add $\lceil \frac{1}{2}(\Delta - 3k) \rceil$ new vertices, each adjacent to v and w . Since H is chordal and each new vertex is simplicial, G is chordal. The addition of degree-2 vertices to H does not increase the maximum clique size (since $k \geq 2$). Thus G has clique number $2k$ and tree-width $2k - 1$. Since each vertex of H is incident to at most two horizontal edges, G has maximum degree $3k - 1 + 2\lceil \frac{1}{2}(\Delta - 3k) \rceil \leq \Delta$.

Observe that $V(G) - V(H)$ is an independent set of simplicial vertices in G . By Lemma 3, G has a tree-partition T with width $\text{tpw}(G)$, such that for every bag B of T , either $B = \{v\}$

FIGURE 2. The graph G with $k = 4$, $\Delta = 15$, and $n = 9$.

for some vertex v of $G - H$, or the induced subgraph $H[B]$ is connected. Since G is connected, T is a (connected) tree. Let U be the tree-partition of H induced by T . That is, to obtain U from T delete the vertices of $G - H$ from each bag, and delete empty bags. Since H is connected, U is a (connected) tree. By Lemma 3, each bag of U induces a connected subgraph of H .

Suppose that U only has two bags B and C . Then one of B and C contains at least $\frac{1}{2}nk$ vertices. Since $k \geq 2$, we have $\text{tpw}(G) \geq \frac{1}{2}nk > \frac{1}{4}k(\Delta - 3k)$, as desired. Now assume that U has at least three bags.

Consider a bag B of U . Let $\ell(B)$ be the minimum integer such that some vertex in B is in the $\ell(B)$ -column, and let $r(B)$ be the maximum integer such that some vertex in B is in the $r(B)$ -column. Since $H[B]$ is connected, there is a path in B from the $\ell(B)$ -column to the $r(B)$ -column. By the definition of H , for each $x \in [\ell(B), r(B)]$, the x -column contains a vertex in B . Let $I(B)$ be the closed real interval from $\ell(B) - \frac{1}{2}$ to $r(B) + \frac{1}{2}$. Observe that two bags B and C of U are adjacent if and only if $I(B) \cap I(C) \neq \emptyset$. Thus $\{I(B) : B \text{ is a bag of } U\}$ is an interval representation of the tree U . Every tree that is an interval graph is a caterpillar⁵; see [16] for example. Thus U is a caterpillar.

Let \preceq be the relation on the set of non-leaf bags of U defined by $A \preceq B$ if and only if $\ell(A) \leq \ell(B)$ and $r(A) \leq r(B)$. We claim that \preceq is a total order. It is immediate that \preceq is reflexive and transitive. To prove that \preceq is antisymmetric, suppose on the contrary that $A \preceq B$ and $B \preceq A$ for distinct non-leaf bags A and B . Thus $\ell(A) = \ell(B)$ and $r(A) = r(B)$.

⁵A *caterpillar* is a tree such that deleting the leaves gives a path.

Since U has at least three bags, there is a third bag C that contains a vertex in the $(\ell(A) - 1)$ -column or in the $(r(A) + 1)$ -column. Thus $\{A, B, C\}$ induce a triangle in U , which is the desired contradiction. Hence \preceq is antisymmetric. To prove that \preceq is total, suppose on the contrary that $A \not\preceq B$ and $B \not\preceq A$ for distinct non-leaf bags A and B . Now $A \not\preceq B$ implies that $\ell(A) > \ell(B)$ or $r(A) > r(B)$. Without loss of generality, $\ell(A) > \ell(B)$. Thus $B \not\preceq A$ implies that $r(B) > r(A)$. Hence the interval $[\ell(A), r(A)]$ is strictly within the interval $[\ell(B), r(B)]$ at both ends. For each $x \in [\ell(A), r(A)]$, every vertex in the x -column is in $A \cup B$, as otherwise U would contain a triangle (since each column is a clique in H). Moreover, every vertex in the $(\ell(A) - 1)$ -column or in the $(r(A) + 1)$ -column is in B , as otherwise U would contain a triangle (since the union of consecutive columns is a clique in H). Thus every neighbour of every vertex in A is in B . That is, A is a leaf in U . This contradiction proves that \preceq is a total order on the set of non-leaf bags of U .

Suppose that U has a 4-vertex path (A, B, C, D) as a subgraph.

Thus B and C are non-leaf bags. Without loss of generality, $B \prec C$. If every column contains vertices in both B and C , then B and C and any other bag would induce a triangle in U (since each column induces a clique in H). Thus some column contains a vertex in B but no vertex in C , and some column contains a vertex in C but no vertex in B . Let p be the maximum integer such that some vertex in B is in the p -column, but no vertex in C is in the p -column. Let q be the minimum integer such that some vertex in C is in the q -column, but no vertex in B is in the q -column. Now $p < q$ since $B \prec C$.

We claim that the $(p + 1)$ -column contains a vertex in C . If not, then the $(p + 1)$ -column contains no vertex in B by the definition of p . Thus $r(B) = p$ since $H[B]$ is connected. Since B is adjacent to C in U , $\ell(C) \leq r(B) + 1 = p + 1$. In particular, the $(p + 1)$ -column contains a vertex in C . Since $H[C]$ is connected, for $x \in [p + 1, q]$, each x -column contains a vertex in C . In fact, $\ell(C) = p + 1$ since the p -column contains no vertex in C . By symmetry, for $x \in [p, q - 1]$, each x -column contains a vertex in B , and $r(C) = q - 1$.

The union of the p -column and the $(p + 1)$ -column only contains vertices in $B \cup C$, as otherwise U would contain a triangle (since the union of two consecutive columns is a clique in H). By the definition of p , no vertex in the p -column is in C . Thus every vertex in the p -column is in B . By symmetry, every vertex in the q -column is in C . Now for each $y \in [k]$, the vertices $(p, y), (p + 1, y), \dots, (q, y)$ are all in $B \cup C$, the first vertex (p, y) is in B , and the last vertex (q, y) is in C . Thus $(x, y) \in B$ and $(x + 1, y) \in C$ for some $x \in [p, q - 1]$. That is, in every row of H there is a horizontal edge with one endpoint in B and the other in C .

Thus there are at least k horizontal edges with one endpoint in B and the other in C (now considered to be bags of T). For each such horizontal edge vw , each vertex of $G - H$ adjacent to v and w is in $B \cup C$, as otherwise T would contain a triangle. There are $\lceil \frac{1}{2}(\Delta - 3k) \rceil$ such vertices of $G - H$ for each of the k horizontal edges between B and C . Thus $|B \cup C| \geq \frac{1}{2}k(\Delta - 3k)$. Thus one of B and C has at least $\frac{1}{4}k(\Delta - 3k)$ vertices. Hence $\text{tpw}(G) \geq \frac{1}{4}k(\Delta - 3k)$ as desired.

Now assume that U has no 4-vertex path as a subgraph.

A tree is a star if and only if it has no 4-vertex path as a subgraph. Hence U is a star. Let R be the root bag of U . If R contains a vertex in every column then $|R| \geq n$, implying

$\text{tpw}(G) \geq n \geq \frac{1}{4}k(\Delta - 3k)$, as desired. Now assume that for some $x \in [n]$, the x -column of H contains no vertex in R . Let B be a bag containing some vertex in the x -column. The x -column induces a clique in H , the only bag in U that is adjacent to B is R , and R contains no vertex in the x -column. Thus every vertex in the x -column is in B . Since R is the only bag in U adjacent to B , there are at least k horizontal edges with one endpoint in B and the other endpoint in R . As in the case when U contained a 4-vertex path, we conclude that $\text{tpw}(G) \geq \frac{1}{4}k(\Delta - 3k)$ as desired. \square

Proof of Theorem 2. Let $\ell := \lceil \frac{k}{2} \rceil$. Thus $\ell \geq 2$. By Lemma 4, for each integer $\Delta \geq \Delta(k, \epsilon) := \max\{3\ell + 1, \frac{3\ell}{8\epsilon}\}$, there are infinitely many values of N for which there is a chordal graph G with N vertices, tree-width $\text{tw}(G) = 2\ell - 1 \leq k$, maximum degree $\Delta(G) \leq \Delta$, and tree-partition-width $\text{tpw}(G) > \frac{1}{4}\ell(\Delta - 3\ell)$, which is at least $(\frac{1}{8} - \epsilon)k\Delta$ since $\Delta \geq \frac{3\ell}{8\epsilon}$. \square

A *domino tree decomposition*⁶ is a tree decomposition in which each vertex appears in at most two bags. The *domino tree-width* of a graph G , denoted by $\text{dtw}(G)$, is the minimum width of a domino tree decomposition of G . Domino tree-width behaves like tree-partition-width in the sense that $\text{dtw}(G) \geq \text{tw}(G)$, and $\text{dtw}(G)$ is bounded for graphs of bounded tree-width and bounded degree [7]. The best upper bound is

$$\text{dtw}(G) \leq (9\text{tw}(G) + 7) \Delta(G) (\Delta(G) + 1) - 1,$$

which is due to Bodlaender [6], who also constructed a graph G with

$$\text{dtw}(G) \geq \frac{1}{12} \text{tw}(G) \Delta(G) - 2.$$

Tree-partition-width and domino tree-width are related in that every graph G satisfies

$$\text{dtw}(G) \geq \text{tpw}(G) - 1,$$

as observed by Bodlaender and Engelfriet [7]. Thus Theorem 2 provides examples of graphs G with

$$\text{dtw}(G) \geq (\frac{1}{8} - \epsilon) \text{tw}(G) \Delta(G).$$

This represents a small constant-factor improvement over the above lower bound by Bodlaender [6].

4. LOWER BOUND FOR TREE-WIDTH 2

We now prove a lower bound on the tree-partition-width of graphs with tree-width 2.

Theorem 3. *For all odd $\Delta \geq 11$ there is a chordal graph G with tree-width 2, maximum degree Δ , and tree-partition-width $\text{tpw}(G) \geq \frac{2}{3}(\Delta - 1)$.*

Proof. As illustrated in Figure 3, let G be the graph with

$$V(G) := \{r\} \cup \{v_i : i \in [\Delta]\} \cup \{w_{i,\ell} : i \in [\Delta - 1], \ell \in [\frac{1}{2}(\Delta - 3)]\}$$

and

$$E(G) := \{rv_i : i \in [\Delta]\} \cup \{v_i v_{i+1} : i \in [\Delta - 1]\} \cup \{v_i w_{i,\ell}, v_{i+1} w_{i,\ell} : i \in [\Delta - 1], \ell \in [\frac{1}{2}(\Delta - 3)]\}.$$

⁶See [10] for an introduction to tree decompositions.

Observe that G has maximum degree Δ . Clearly every induced cycle of G is a triangle. Thus G is chordal. Observe that G has no 4-vertex clique. Thus G has tree-width 2.




FIGURE 3. Illustration for Theorem 3 with $\Delta = 13$.

Let T be the tree-partition of G from Lemma 3. Then T has width $\text{tpw}(G)$, and every bag induces a connected subgraph of G . Let R be the bag containing r . Let B_1, \dots, B_d be the bags, not including R , that contain some vertex v_i . Thus R is adjacent to each B_j (since r is adjacent to each v_i). Since $\{w_{i,\ell} : i \in [\Delta - 1], \ell \in [\frac{1}{2}(\Delta - 3)]\}$ is an independent set of simplicial vertices, by Lemma 3, for each $j \in [d]$, the vertices $\{v_1, v_2, \dots, v_\Delta\} \cap B_j$ induce a (connected) subpath of G .

First suppose that $d = 0$. Then the $\Delta + 1$ vertices $\{r, v_1, \dots, v_\Delta\}$ are contained in one bag R . Thus $\text{tpw}(G) \geq \Delta + 1 \geq \frac{2}{3}(\Delta - 1)$.

Now suppose that $d = 1$. Thus $\{r, v_1, \dots, v_\Delta\} \subseteq R \cup B_1$. In addition, at least one edge $v_i v_{i+1}$ has one endpoint in R and the other endpoint in B_1 . Thus $w_{i,\ell} \in R \cup B_1$ for each $\ell \in [\frac{1}{2}(\Delta - 3)]$. Hence $1 + \Delta + \frac{1}{2}(\Delta - 3)$ vertices are contained in two bags. Thus one bag contains at least $\frac{1}{4}(3\Delta - 1)$ vertices, and $\text{tpw}(G) \geq \frac{1}{4}(3\Delta - 1) \geq \frac{2}{3}(\Delta - 1)$.

Finally suppose that $d \geq 2$. Since $\{v_1, v_2, \dots, v_\Delta\} \cap B_j$ induce a subpath in each bag B_j , we can assume that $\{v_1, v_2, \dots, v_\Delta\} \cap B_j = \{v_i : i \in [f(j), g(j)]\}$, where

$$1 \leq f(1) \leq g(1) < f(2) \leq g(2) < \dots < f(d) \leq g(d) \leq \Delta.$$

Distinct B_j bags are not adjacent (since T is a tree). Thus $v_{f(j)-1} \in R$ for each $j \in [2, d]$. Similarly, $v_{g(j)+1} \in R$ for each $j \in [d - 1]$. Thus $w_{f(j)-1,\ell} \in R \cup B_j$ for each $j \in [2, d]$ and $\ell \in [\frac{1}{2}(\Delta - 3)]$. Similarly, $w_{g(j),\ell} \in R \cup B_j$ for each $j \in [d - 1]$ and $\ell \in [\frac{1}{2}(\Delta - 3)]$.

Hence the bags R, B_1, \dots, B_d contain at least

$$1 + \Delta + 2(d - 1) \cdot \frac{1}{2}(\Delta - 3)$$


vertices. Therefore one of these bags has at least

$$(1 + \Delta + (d - 1)(\Delta - 3)) / (d + 1)$$

vertices, which is at least $\frac{2}{3}(\Delta - 1)$. Hence $\text{tpw}(G) \geq \frac{2}{3}(\Delta - 1)$. \square

REFERENCES

- [1] NOGA ALON, GUOLI DING, BOGDAN OPOROWSKI, AND DIRK VERTIGAN. Partitioning into graphs with only small components. *J. Combin. Theory Ser. B*, 87(2):231–243, 2003.
- [2] JÁNOS BARÁT AND DAVID R. WOOD. Notes on nonrepetitive graph colouring, 2005. [arXiv.org/math/0509608](https://arxiv.org/math/0509608).

FIGURE 4. Illustration for Theorem 3 with $\Delta = 19$ and $d = 4$.

- [3] HANS L. BODLAENDER. The complexity of finding uniform emulations on fixed graphs. *Inform. Process. Lett.*, 29(3):137–141, 1988.
- [4] HANS L. BODLAENDER. The complexity of finding uniform emulations on paths and ring networks. *Inform. and Comput.*, 86(1):87–106, 1990.
- [5] HANS L. BODLAENDER. A partial k -arboretum of graphs with bounded treewidth. *Theoret. Comput. Sci.*, 209(1-2):1–45, 1998.
- [6] HANS L. BODLAENDER. A note on domino treewidth. *Discrete Math. Theor. Comput. Sci.*, 3(4):141–150, 1999.
- [7] HANS L. BODLAENDER AND JOOST ENGELFRIET. Domino treewidth. *J. Algorithms*, 24(1):94–123, 1997.
- [8] HANS L. BODLAENDER AND JAN VAN LEEUWEN. Simulation of large networks on smaller networks. *Inform. and Control*, 71(3):143–180, 1986.
- [9] EMILIO DI GIACOMO, GIUSEPPE LIOTTA, AND HENK MEIJER. Computing straight-line 3D grid drawings of graphs in linear volume. *Comput. Geom.*, 32(1):26–58, 2005.
- [10] REINHARD DIESTEL. *Graph theory*, vol. 173 of *Graduate Texts in Mathematics*. Springer, 2nd edn., 2000. ISBN 0-387-95014-1.
- [11] REINHARD DIESTEL AND DANIELA KÜHN. Graph minor hierarchies. *Discrete Appl. Math.*, 145(2):167–182, 2005.
- [12] GUOLI DING AND BOGDAN OPOROWSKI. Some results on tree decomposition of graphs. *J. Graph Theory*, 20(4):481–499, 1995.
- [13] GUOLI DING AND BOGDAN OPOROWSKI. On tree-partitions of graphs. *Discrete Math.*, 149(1-3):45–58, 1996.
- [14] VIDA DUJMOVIĆ, PAT MORIN, AND DAVID R. WOOD. Layout of graphs with bounded tree-width. *SIAM J. Comput.*, 34(3):553–579, 2005.
- [15] VIDA DUJMOVIĆ, MATTHEW SUDERMAN, AND DAVID R. WOOD. Graph drawings with few slopes. *Comput. Geom.*, 38:181–193, 2007.
- [16] JÜRGEN ECKHOFF. Extremal interval graphs. *J. Graph Theory*, 17(1):117–127, 1993.
- [17] ANDERS EDENBRANDT. Quotient tree partitioning of undirected graphs. *BIT*, 26(2):148–155, 1986.

- [18] JOHN P. FISHBURN AND RAPHAEL A. FINKEL. Quotient networks. *IEEE Trans. Comput.*, C-31(4):288–295, 1982.
- [19] RUDOLF HALIN. Tree-partitions of infinite graphs. *Discrete Math.*, 97:203–217, 1991.
- [20] DIETRICH KUSKE AND MARKUS LOHREY. Logical aspects of Cayley-graphs: the group case. *Ann. Pure Appl. Logic*, 131(1–3):263–286, 2005.
- [21] BRUCE A. REED. Algorithmic aspects of tree width. In BRUCE A. REED AND CLÁUDIA L. SALES, eds., *Recent Advances in Algorithms and Combinatorics*, pp. 85–107. Springer, 2003.
- [22] NEIL ROBERTSON AND PAUL D. SEYMOUR. Graph minors. II. Algorithmic aspects of tree-width. *J. Algorithms*, 7(3):309–322, 1986.
- [23] DETLEF SEESE. Tree-partite graphs and the complexity of algorithms. In LOTHAR BUDACH, ed., *Proc. International Conf. on Fundamentals of Computation Theory*, vol. 199 of *Lecture Notes in Comput. Sci.*, pp. 412–421. Springer, 1985.
- [24] DAVID R. WOOD. Vertex partitions of chordal graphs. *J. Graph Theory*, 53(2):167–172, 2006.
- [25] DAVID R. WOOD AND JAN ARNE TELLE. Planar decompositions and the crossing number of graphs with an excluded minor. *New York J. Math.*, 13:117–146, 2007.

DEPARTAMENT DE MATEMÀTICA APLICADA II
 UNIVERSITAT POLITÈCNICA DE CATALUNYA
 BARCELONA, SPAIN

E-mail address: david.wood@upc.es